




PULSION 社 PiCCO 関連文献リスト



TOKIBO
CO., LTD.

PULSION PiCCO 関連文献リスト

目次

1. レビュー文献 (REVIEWS)	3
1.1 一般 (GENERAL)	3
1.2 心拍出量 (CARDIAC OUTPUT)	4
1.3 前負荷 (PRELOAD)	4
1.4 肺血管外水分量 (LUNG WATER)	4
1.5 小児 (PAEDIATRIC)	5
2. ガイドラインやスタンダードな手技 (GUIDELINES AND STANDARD OPERATING PROCEDURES)	5
3a. パラメータの理論 (PICCO PARAMETERS METHODOLOGY)	6
3a.1 心拍出量 (FLOW (Cardiac Output))	6
3a.2 前負荷 (PRELOAD (GEDV and ITBV))	6
3a.3 収縮性 (CONTRACTILITY (GEF, CFI & Left Ventricular Contractility dPmx))	7
3a.4 輸液反応 (VOLUME RESPONSIVENESS (SVV and PPV))	7
3a.5 肺水腫 (PULMONARY EDEMA (EVLW))	7
3b. パラメータの検証 (PICCO PARAMETERS VALIDATION)	8
3b.1 心拍出量 (FLOW (Cardiac Output CO))	8
3b.2 前負荷 (PRELOAD (GEDV and ITBV))	9
3b.3 収縮性 (CONTRACTILITY (GEF, CFI & Left Ventricular Contractility dPmx))	9
3b.4 輸液反応 (VOLUME RESPONSIVENESS (SVV and PPV))	10
3b.5 肺水腫 (PULMONARY EDEMA (EVLW))	10
4. 適用範囲 (FIELDS OF APPLICATION)	11
4.1 メディカル ICU (MEDICAL INTENSIVE CARE)	11
4.2 サージカル ICU (SURGICAL INTENSIVE CARE)	11
4.3 心臓手術 (CARDIAC SURGERY)	12
4.4 移植 (TRANSPLANTATION)	14
4.5 外傷 (TRAUMA)	14
4.6 熱傷 (BURNS)	15
4.7 小児 (PEDIATRIC)	15
4.8 敗血症性ショック (SHOCK SEPSIS)	16
4.9 出血性ショック (SHOCK HYPOVOLEMIA)	17
4.10 心原性ショック (SHOCK CARDIOGENIC)	13
4.11 ARDS と ALI (ACUTE RESPIRATORY DISTRESS SYNDROME / ACUTE LUNG INJURY)	18
4.12 その他 (OTHERS)	19
脳神経外科関連 (NEUROSURGERY / NEUROLOGY)	19
右-左シャント (RIGHT-LEFT SHUNT)	19
肺水腫 (PULMONARY EDEMA)	19
5. 各パラメータによる治療補助 (DECISION SUPPORT BY PICCO PARAMETERS)	20
6. 様々な臨床状態におけるパラメータ (PICCO PARAMETERS IN VARIOUS CLINICAL SITUATIONS)	21
7. カテーテル関連 (CATHETERS)	22

PULSION PiCCO 文献リスト 推奨度指標

★★★ 非常に強い推奨文献 (Very Highly Recommended)

★★ 強い推奨文献 (Highly Recommended)

★ 推奨文献 (Recommended)

1. レビュー文献 (REVIEWS)

1.1 一般 (GENERAL)

[Malbrain M, De Potter P, Deeren D.](#)

[Cost Effectiveness of minimally invasive hemodynamic monitoring](#)

[In: Vincent JL \(Ed.\), Yearbook of Intensive Care and Emergency Medicine 2005: 603-18](#)

★★★

[Stubbe H, Schmidt C, Hinder F.](#)

[\[Invasives Kreislaufmonitoring: Vier Methoden im Vergleich\] \(Article in German\)](#)

[Anesthesiol Intensivmed Notfallmed Schmerzther 2006; 41\(9\):550-4](#)

★★

[Della Rocca G, Costa MG.](#)

[Volumetric monitoring: principles of application](#)

[Minerva Anesthesiol 2005; 71:303-6](#)

★★★

[Genahr A, McLuckie A.](#)

[Transpulmonary thermodilution in the critically ill](#)

[Brit J Int Care 2004: 6-10](#)

★★★

[Cottis R, Magee N, Higgins DJ.](#)

[Haemodynamic monitoring with pulse-induced contour cardiac output \(PiCCO\) in critical care](#)

[Intensive Crit Care Nurs 2003; 19: 301-7](#)

★★

[Michard F, Perel A.](#)

[Management of circulatory and respiratory failure using less invasive hemodynamic monitoring](#)

[In: Vincent JL \(Ed.\), Yearbook of Intensive Care and Emergency Medicine 2003: 508-20](#)

★★★

[Bellomo R, Uchino S.](#)

[Cardiovascular monitoring tools: use and misuse](#)

[Curr Opin Crit Care 2003; 9\(3\): 225-9](#)

★★

1.2 心拍出量(CARDIAC OUTPUT)

[Reuter D, Goetz AE.](#)

[\[Messung des Herzzeitvolumens\] \(Article in German\)](#)

[Anaesthesist 2005; 54:1135–53](#)

[★★★](#)

1.3 前負荷(PRELOAD)

[Della Rocca G, Costa MG, Pietropaoli P.](#)

[How to measure and interpret volumetric measures of preload](#)

[Curr Opin Crit Care 2007; 13\(3\): 297–302](#)

[★★](#)

[Della Rocca G, Costa MG.](#)

[Intrathoracic Blood Volume: Clinical Applications](#)

[In: Vincent JL \(Ed.\), Yearbook of Intensive Care and Emergency Medicine 2006: 143–52](#)

[★★★](#)

[Michard F.](#)

[Do we need to know cardiac preload?](#)

[In: Vincent JL \(Ed.\), Yearbook of Intensive Care and Emergency Medicine 2004: 694–701](#)

[★★★](#)

1.4 肺血管外水分量(LUNG WATER)

[Groeneveld ABJ.](#)

[Value and limitations of measuring extravascular lung water](#)

[Int J Intensive Care 2007; Autumn: 84–6](#)

[★](#)

[Khan S, Trof RJ, Groeneveld ABJ.](#)

[Transpulmonary dilution-derived extravascular lung water as a measure of lung edema](#)

[Curr Opin Crit Care 2007; 13 \(3\):303–7](#)

[★★★](#)

[Fernandez-Mondejar E, Guerrero-Lopez F, Colmenero M.](#)

[How important is the measurement of extravascular lung water?](#)

[Curr Opin Crit Care 2007; 13: 79–83](#)

[★★★](#)

[Michard F.](#)

[Bedside assessment of extravascular lung water by dilution methods: temptations and pitfalls](#)

[Crit Care Med 2007; 35\(4\):1186–92](#)

[★★★](#)

[Isakow W, Schuster DP.](#)

[Extravascular lung water measurements and hemodynamic monitoring in the critically ill: bedside alternatives to the pulmonary artery catheter](#)

[Am J Physiol Lung Cell Mol Physiol 2006; 291: 1118–33](#)

★★★

[Kirov MY, Kuzkov VV, Bjertnaes LJ.](#)

[Extravascular lung water in sepsis](#)

[In: Vincent JL \(Ed.\), Yearbook of Intensive Care and Emergency Medicine 2005: 449–60](#)

★★★

1.5 小児 (PAEDIATRIC)

[Tibby SM, Murdoch IA.](#)

[Measurement of cardiac output and tissue perfusion](#)

[Curr Opin Pediatr 2002; 14\(3\):303–9](#)

★★

2. ガイドラインやスタンダードな手技 (GUIDELINES AND STANDARD OPERATING PROCEDURES)

[Goepfert M, Reuter D, Akyol D, Lamm P, Kilger E, Goetz A.](#)

[Goal directed fluid management reduces vasopressor and catecholamine use in cardiac surgery patients](#)

[Intensive Care Medicine 2007; 33: 96–103](#)

★★★

[Kortgen A, Niederprun P, Bauer M.](#)

[Implementation of an evidence-based “standard operating procedure” and outcome in septic shock](#)

[Crit Care Med 2006; 34\(4\):939–9](#)

★★★

[Reinhart K, Brunkhorst FM, Bone HG, Gerlach H, Grundling M, Kreymann G, Kujath P, Marggraf G, Mayer K, Meier-Hellmann A, Peckelsen C, Putensen C, Quintel M, Ragaller M, Rossaint R, Stuber F, Weiler N, Welte T, Werdan K.](#)

[\[Diagnose und Therapie der Sepsis: S2-Leitlinien der Deutschen Sepsis-Gesellschaft e.V. \(DSG\) und der Deutschen Interdisziplinären Vereinigung für Intensiv- und Notfallmedizin \(DIVI\)\] \(Article in German\)](#)

[Internist 2006:1–40](#)

★★★

[Hensel M, Schwenk W, Bloch A, Raue W, Stracke S, Volk T, v. Heymann C, Müller JM, Kox WJ, Spies C.](#)

[\[Die Aufgabe der Anesthesiologie bei der Umsetzung operativer „Fast track-Konzepte“\] \(Article in German\)](#)

[Der Anesthesist 2006; 5\(1\):80–92](#)

★★

[Dellinger RP, Carlet JM, Masur H, Gerlach H, Calandra T, Cohen J, Gea-Banacloche J, Keh D, Marshall JC, Parker MM, Ramsay G, Zimmerman JL, Vincent JL, Levy MM. Surviving Sepsis Campaign guidelines for management of severe sepsis and septic shock Crit Care Med 2004; 32\(3\): 858-77](#)

★★

3a. パラメータの理論 (PICCO PARAMETERS METHODOLOGY)

3a.1 心拍出量 (FLOW (Cardiac Output))

[Faybik P, Hetz H, Baker A, Yankovskaya E, Krenn CG, Steltzer H. Iced versus room temperature injectate for assessment of cardiac output, intrathoracic blood volume, and extravascular lung water by single transpulmonary thermodilution J Crit Care 2004; 19\(2\):103-7](#)

★

[Segal E, Katzenelson R, Berkenstadt H, Perel A. Transpulmonary thermodilution cardiac output measurement using the axillary artery in critically ill patients J Clin Anesth 2002; 14\(3\):210-3](#)

★★★

[Sakka SG, Meier-Hellmann A. Evaluation of cardiac output and cardiac preload In: Vincent JL \(Ed.\), Yearbook of Intensive Care and Emergency Medicine 2000: 671-9](#)

★★

3a.2 前負荷 (PRELOAD (GEDV and ITBV))

[Hofer CK, Furrer L, Matter-Ensner S, Maloigne M, Klaghofer R, Genoni M, Zollinger A. Volumetric preload measurement by thermodilution: a comparison with transoesophageal echocardiography Br J Anaesth 2005; 94\(6\):748-55](#)

★★★

[Buhre W, Buhre K, Kazmaier S, Sonntag H, Weyland A. Assessment of cardiac preload by indicator dilution and transoesophageal echocardiography Eur J Anaesthesiol 2001; 18\(10\):662-7](#)

★★★

[McLuckie A, Bihari D. Investigating the relationship between intrathoracic blood volume index and cardiac index Intensive Care Med 2000; 26\(9\):1376-8](#)

★★★

3a.3 收縮性(CONTRACTILITY (GEF, CFI & Left Ventricular Contractility dPmx))

[Michard F, Perel A.](#)

[Management of circulatory and respiratory failure using less invasive hemodynamic monitoring](#)

[In: Vincent JL \(Ed.\), Yearbook of Intensive Care and Emergency Medicine 2003: 508-20](#)

★★★

[Wisner-Euteneier AJ, Lichtwarck-Aschoff M, Zimmermann G, Bluemel G, Pfeiffer U.](#)

[Evaluation of the cardiac function index as a new bedside indicator of cardiac performance](#)

[Intensive Care Med 1994; 20\(S2\):21](#)

★★

3a.4 輸液反応(VOLUME RESPONSIVENESS (SVV and PPV))

[Reuter DA, Goepfert MS, Goresch T, Schmoeckel M, Kilger E, Goetz AE.](#)

[Assessing fluid responsiveness during open chest conditions](#)

[Br J Anaesth 2005; 94\(3\):318-23](#)

★★★

[Rex S, Brose S, Metzelder S, Huneke R, Schalte G, Autschbach R, Rossaint R, Buhre W.](#)

[Prediction of fluid responsiveness in patients during cardiac surgery](#)

[Br J Anaesth 2004; 93 \(6\):782-8](#)

★★★

[Reuter DA, Felbinger TW, Kilger E, Schmidt C, Lamm P, Goetz AE.](#)

[Optimizing fluid therapy in mechanically ventilated patients after cardiac surgery by on-line monitoring of left ventricular stroke volume variations. Comparison with aortic systolic pressure variations](#)

[Br J Anaesth 2002; 88\(1\):124-6](#)

★★★

3a.5 肺水腫(PULMONARY EDEMA (Extravascular Lung Water))

[Phillips, C., Chesnutt M, Smith, M.](#)

[Extravascular lung water in sepsis-associated acute respiratory distress syndrome: indexing with predicted body weight improves correlation with severity of illness and survival](#)

[Crit Care Med, 2007\(epub\): 1-5](#)

★★★

[Isakow W, Schuster DP.](#)

[Extravascular lung water measurements and hemodynamic monitoring in the critically ill: bedside alternatives to the pulmonary artery catheter](#)

[Am J Physiol Lung Cell Mol Physiol 2006: 291\(6\):1118-33](#)

★★★

[Kuzkov V, Kirov M, Sovershaev M, Kulkin V, Suberov E, Waerhaug K, Bjertnaes L.
Extravascular lung water determined with single transpulmonary thermodilution correlates with the
severity of sepsis-induced acute lung injury
Crit Care Med 2006; 34\(6\):1647-53](#)

★★★

[Kirov MY, Kuzkov VV, Bjertnaes LJ.
Extravascular lung water in sepsis
In: Vincent JL \(Ed.\), Yearbook of Intensive Care and Emergency Medicine 2005: 449-60](#)

★★★

[Sakka SG, Klein M, Reinhart K, Meier-Hellmann A.
Prognostic value of extravascular lung water in critically ill patients
Chest 2002; 122\(6\):2080-6](#)

★★★

3b. パラメータの検証 (PICCO PARAMETERS VALIDATION)

3b.1 心拍出量 (FLOW (Cardiac Output CO))

[Felbinger TW, Reuter DA, Eltzschig HK, Bayerlein J, Goetz AE.
Cardiac index measurements during rapid preload changes: a comparison of pulmonary artery
thermodilution with arterial pulse contour analysis
J Clin Anesth 2005; 17\(4\):241-8](#)

★★★

[Marx G, Schuerholz T, Sumpelmann R, Simon T, Leuwer M.
Comparison of cardiac output measurements by arterial trans-cardiopulmonary and pulmonary arterial
thermodilution with direct Fick in septic shock
Eur J Anaesthesiol 2005; 22\(2\):129-34](#)

★★★

[Bein B, Worthmann F, Tonner PH, Paris A, Steinfath M, Hedderich J, Scholz J.
Comparison of esophageal Doppler, pulse contour analysis, and real-time pulmonary artery
thermodilution for the continuous measurement of cardiac output
J Cardiothorac Vasc Anesth 2004; 18\(2\):185-9](#)

★★★

[Godje O, Hoke K, Goetz AE, Felbinger TW, Reuter DA, Reichart B, Friedl R, Hannekum A, Pfeiffer UJ.
Reliability of a new algorithm for continuous cardiac output determination by pulse-contour analysis
during hemodynamic instability
Crit Care Med 2002; 30\(1\):52-8](#)

★★★

[Sakka SG, Reinhart K, Wegscheider K, Meier-Hellmann A.](#)

[Is the placement of a pulmonary artery catheter still justified solely for the measurement of cardiac output?](#)

[J Cardiothorac Vasc Anesth 2000; 14\(2\):119-24](#)

★★★

3b.2 前負荷 (PRELOAD (GEDV and ITBV))

[Renner J, Gruenewald M, Brand P, Steinfath M, Scholz J, Lutter G, Bein B.](#)

[Global End-Diastolic Volume as a Variable of Fluid Responsiveness During Acute Changing Loading Conditions](#)

[J Cardiothorac Vasc Anesth 2007; 21\(5\): 650-4](#)

★★

[Kozieras, J, Thuemer O, Sakka SG.](#)

[Influence of an acute increase in systemic vascular resistance on transpulmonary thermodilution-derived parameters in critically ill patients](#)

[Intensive Care Med 2007; 33:1619-23](#)

★★★

[Michard F, Alaya S, Zarka V, Bahloul M, Richard C, Teboul JL.](#)

[Global end-diastolic volume as an indicator of cardiac preload in patients with septic shock](#)

[Chest 2003; 124\(5\):1900-8](#)

★★★

[Reuter DA, Felbinger TW, Moerstedt K, Weis F, Schmidt C, Kilger E, Goetz AE.](#)

[Intrathoracic blood volume index measured by thermodilution for preload monitoring after cardiac surgery](#)

[J Cardiothorac Vasc Anesth 2002; 16\(2\):191-5](#)

★★★

3b.3 収縮性 (CONTRACTILITY (GEF, CFI & Left Ventricular Contractility dPmx))

[de Hert S, Robert D, Cromheecke S, Michard F, Nijs J, Rodrigues IE.](#)

[Evaluation of Left Ventricular Function in Anesthetised Patients Using Femoral Artery dP/dtmax](#)

[J Cardio Thor Vasc Anes 2006; 20\(3\): 325-30](#)

★★★

[Combes A, Berneau JB, Luyt CE, Trouillet JL.](#)

[Estimation of left ventricular systolic function by single transpulmonary thermodilution](#)

[Intensive Care Med 2004; 30\(7\):1377-83](#)

★★

3b.4 輸液反応(VOLUME RESPONSIVENESS (SVV and PPV))

[Kubitz JC, Anneck T, Forkl S, Kemming GI, Kronas N, Goetz AE, Reuter DA.](#)

[Validation of pulse contour derived stroke volume variation during modifications of cardiac afterload](#)

[Br J Anaesth 2007; 98\(5\): 591–7](#)

★★

[Hofer CK, Muller SM, Furrer L, Klaghofer R, Genoni M, Zollinger A.](#)

[Stroke volume and pulse pressure variation for prediction of fluid responsiveness in patients undergoing off-pump coronary artery bypass grafting](#)

[Chest 2005; 128\(2\):848–54](#)

★★★

[Reuter DA, Kirchner A, Felbinger TW, Weis FC, Kilger E, Lamm P, Goetz AE.](#)

[Usefulness of left ventricular stroke volume variation to assess fluid responsiveness in patients with reduced cardiac function](#)

[Crit Care Med 2003; 31\(5\):1399–404](#)

★★★

[Michard F, Boussat S, Chemla D, Anguel N, Mercat A, Lecarpentier Y, Richard C, Pinsky MR, Teboul JL.](#)

[Relation between respiratory changes in arterial pulse pressure and fluid responsiveness in septic patients with acute circulatory failure](#)

[Am J Respir Crit Care Med 2000; 162\(1\):134–8](#)

★★

3b.5 肺水腫(PULMONARY EDEMA (EVLW))

[Monnet X, Anguel N, Osman D, Hamzaoui, Richard C, Teboul JL.](#)

[Assessing pulmonary permeability by transpulmonary thermodilution allows differentiation of hydrostatic pulmonary edema from ALI / ARDS](#)

[Intensive Care Medicine 2007; 33\(3\):448–53](#)

★★★

[Kuzkov VV, Suborov EV, Kirov MY, Kuklin VN, Sobhkezh M, Johnsen S, Waerhaug K, Bjertnaes LJ.](#)

[Extravascular lung water after pneumonectomy and one-lung ventilation in sheep](#)

[Crit Care Med 2007; 35 \(6\):1550–9](#)

★★★

[Michard F, Schachtrupp A, Toens C.](#)

[Factors influencing the estimation of extravascular lung water by transpulmonary thermodilution in critically ill patients](#)

[Crit Care Med 2005; 33\(6\):1243–7](#)

★★★

[Kirov MY, Kuzkov VV, Kuklin VN, Waerhaug K, Bjertnaes LJ.](#)

[Extravascular lung water assessed by transpulmonary single thermodilution and postmortem gravimetry in sheep](#)

[Crit Care 2004; 8\(6\):R451-8](#)

★★★

[Katzenelson R, Perel A, Berkenstadt H, Preisman S, Kogan S, Sternik L, Segal E.](#)

[Accuracy of transpulmonary thermodilution versus gravimetric measurement of extravascular lung water](#)

[Crit Care Med 2004; 32\(7\):1550-4](#)

★★★

[Sakka SG, Ruhl CC, Pfeiffer UJ, Beale R, McLuckie A, Reinhart K, Meier-Hellmann A.](#)

[Assessment of cardiac preload and extravascular lung water by single transpulmonary thermodilution](#)

[Intensive Care Medicine 2000; 26 \(2\):180-7](#)

★★★

4. 適用範囲 (FIELDS OF APPLICATION)

4.1 メディカル ICU (MEDICAL INTENSIVE CARE)

[Kortgen A, Niederprun P, Bauer M.](#)

[Implementation of an evidence-based "standard operating procedure" and outcome in septic shock](#)

[Crit Care Med 2006; 34 \(4\):939-9](#)

★★★

[Martin GS, Eaton S, Mealer M, Moss M.](#)

[Extravascular lung water in patients with severe sepsis: a prospective cohort study](#)

[Crit Care 2005; 9\(2\):R74-82](#)

★★★

[Mitchell JP, Schuller D, Calandrino FS, Schuster DP.](#)

[Improved outcome based on fluid management in critically ill patients requiring pulmonary artery catheterization](#)

[Am Rev Respir Dis 1992; 145\(5\):990-8](#)

★★★

4.2 サージカル ICU (SURGICAL INTENSIVE CARE)

[Sato Y, Motoyama S, Maruyama M, Hayashi K, Nakae H, Tajimi K, Ogawa J.](#)

[Extravascular Lung Water Measured Using Single Transpulmonary Thermodilution Reflects Perioperative Pulmonary Edema Induced by Esophagectomy](#)

[Eur Surg Res 2006; 39\(1\): 7-13](#)

★★★

[Roch A, Michelet P, D'Journo B, Brousse D, Blayac D, Lambert D, Auffray JP.](#)

[Accuracy and limits of transpulmonary dilution methods in estimating extravascular lung water after pneumonectomy](#)

[Chest 2005; 128\(2\):927-33](#)

★★

[Deja M, Hildebrandt B, Ahlers O, Riess H, Wust P, Gerlach H, Kerner T.](#)

[Goal-directed therapy of cardiac preload in induced whole-body hyperthermia](#)

[Chest 2005; 128\(2\):580-6](#)

★★

[Michard F, Schachtrupp A, Toens C.](#)

[Factors influencing the estimation of extravascular lung water by transpulmonary thermodilution in critically ill patients](#)

[Crit Care Med 2005; 33\(6\):1243-7](#)

★★★

[Junghans T, Neuss H, Strohauser M, Raue W, Haase O, Schink T, Schwenk W.](#)

[Hypovolemia after traditional preoperative care in patients undergoing colonic surgery is underrepresented in conventional hemodynamic monitoring](#)

[Int J Colorectal Dis 2005; 21\(7\): 693-7](#)

★

[Della Rocca G, Costa, MG, Coccia C, Pompei L, Pietropaoli P.](#)

[Preload and haemodynamic assessment during liver transplantation: a comparison between the pulmonary artery catheter and transpulmonary indicator dilution techniques](#)

[Eur J Anaesthesiol 2002; 19\(12\):868-75](#)

★★★

[Antonini M, Meloncelli S, Dantimi C, Tosti S, Ciotti L, Gasparetto A.](#)

[\[The PiCCO system with brachial-axillary artery access in hemodynamic monitoring during surgery of abdominal aortic aneurysm\] \(Article in Italian\)](#)

[Minerva Anesthesiol 2001; 67\(6\):447-56](#)

★★

4.3 心臟手術 (CARDIAC SURGERY)

[Goepfert M, Reuter D, Akyol D, Lamm P, Kilger E, Goetz A.](#)

[Goal directed fluid management reduces vasopressor and catecholamine use in cardiac surgery patients](#)

[Intensive Care Medicine 2007; 33: 96-103](#)

★★★

[Kirov MY, Lenkin AI, Kuzkov V, Suburov EV, Slastilin VY, Borodin VV, Chernov II, Shonbin AN, Bjertnaes LJ. Single transpulmonary thermodilution in off-pump coronary artery bypass grafting: haemodynamic changes and effects of different anaesthetic techniques](#)
[Acta Anesthesiol Scand 2007; 51: 426-33](#)

★★

[Wouters PF, Quaghebeur B, Sergeant P, Van Hemelrijck J, Vandermeersch E. Cardiac output monitoring using a brachial arterial catheter during off-pump coronary artery bypass grafting](#)
[J Cardiothorac Vasc Anesth 2005; 19: 160-4](#)

★★★

[Reuter DA, Goepfert MS, Goresch T, Schmoeckel M, Kilger E, Goetz AE. Assessing fluid responsiveness during open chest conditions](#)
[Br J Anaesth 2005; 94\(3\):318-23](#)

★★★

[Bettex DA, Hinselmann V, Hellermann JP, Jenni R, Schmid ER. Transoesophageal echocardiography is more unreliable for cardiac output assessment after cardiac surgery compared with thermodilution](#)
[Anesthesia 2004; 59:1184-92](#)

★★

[Mielck F, Buhre W, Hanekop G, Tirilomis T, Hilgers R, Sonntag H. Comparison of continuous cardiac output measurements in patients after cardiac surgery](#)
[J Cardiothorac Vasc Anesth 2003; 17\(2\):211-6](#)

★★★

[Brock H, Gabriel C, Bibl D, Necek S. Monitoring intravascular volumes for postoperative volume therapy](#)
[Eur J Anaesthesiol 2002; 19\(4\):288-94](#)

★★

[Buhre W, Weyland A, Kazmaier S, Hanekop GG, Baryalei MM, Sydow M, Sonntag H. Comparison of cardiac output assessed by pulse-contour analysis and thermodilution in patients undergoing minimally invasive direct coronary artery bypass grafting](#)
[J Cardiothorac Vasc Anesth 1999; 13\(4\):437-40](#)

★★★

[Godje O, Hoke K, Lamm P, Schmitz C, Thiel C, Weinert M, Reichart B. Continuous, less invasive, hemodynamic monitoring in intensive care after cardiac surgery](#)
[Thorac Cardiovasc Surg 1998; 46\(4\):242-9](#)

★★

4.4 移植 (TRANSPLANTATION)

[Della Rocca G, Costa GM, Coccia C, Pompei L, Di Marco P, Pietropaoli P.](#)

[Preload index: pulmonary artery occlusion pressure versus intrathoracic blood volume monitoring during lung transplantation](#)

[Anesth Analg 2002; 95\(4\):835-43](#)

★★★

[Della Rocca G, Costa MG, Coccia C, Pompei L, Pietropaoli P.](#)

[Preload and haemodynamic assessment during liver transplantation: a comparison between the pulmonary artery catheter and transpulmonary indicator dilution techniques](#)

[Eur J Anaesthesiol 2002; 19\(12\):868-75](#)

★★★

[Della Rocca G, Costa MG, Pompei L, Coccia C, Pietropaoli P.](#)

[Continuous and intermittent cardiac output measurement: pulmonary artery catheter versus aortic transpulmonary technique](#)

[Br J Anaesth 2002; 88\(3\):350-6](#)

★★

[Goedje O, Seebauer T, Peyerl M, Pfeiffer UJ, Reichart B.](#)

[Hemodynamic monitoring by double-indicator dilution technique in patients after orthotopic heart transplantation](#)

[Chest 2000; 118\(3\):775-81](#)

★★★

[Krenn CG, Plochl W, Nikolic A, Metnitz PG, Scheuba C, Spiss CK, Steltzer H.](#)

[Intrathoracic fluid volumes and pulmonary function during orthotopic liver transplantation](#)

[Transplantation 2000; 69\(11\):2394-400](#)

★★★

4.5 外傷 (TRAUMA)

[Berkenstadt H, Friedman Z, Preisman S, Keidan I, Livingstone D, Perel A.](#)

[Pulse pressure and stroke volume variations during severe haemorrhage in ventilated dogs](#)

[Br J Anaesth 2005; 94 \(6\): 721-6](#)

★★

[Friedman Z, Berkenstadt H, Margalit N, Segal E, Perel A.](#)

[Cardiac output assessed by arterial thermodilution during exsanguination and fluid resuscitation: experimental validation against a reference technique](#)

[Eur J Anaesthesiol 2002; 19\(5\):337-40](#)

★★

4.6 熱傷 (BURNS)

[Holm C, Mayr M, Horbrand F, Tegeler J, Henckel von Donnersmarck G, Muhlbauer W, Pfeiffer UJ. Reproducibility of transpulmonary thermodilution measurements in patients with burn shock and hypothermia](#)

[J Burn Care Rehabil 2005; 26\(3\):260–5](#)

★★★

[Holm C, Tegeler J, Mayr M, Pfeiffer U, Henckel von Donnersmarck G, Muhlbauer W.](#)

[Effect of crystalloid resuscitation and inhalation injury on extravascular lung water: clinical implications](#)
[Chest 2002; 121\(6\):1956–62](#)

★★

[Holm C, Melcer B, Horbrand F, Henckel von Donnersmarck G, Muhlbauer W.](#)

[Arterial thermodilution: an alternative to pulmonary artery catheter for cardiac output assessment in burn patients](#)

[Burns 2001; 27\(2\):161–6](#)

★★

[Holm C, Melcer B, Horbrand F, Worl H, von Donnersmarck GH, Muhlbauer W.](#)

[Intrathoracic blood volume as an end point in resuscitation of the severely burned: an observational study of 24 patients](#)

[J Trauma 2000; 48\(4\):728–34](#)

★★★

4.7 小児 (PEDIATRIC)

[Fakler U, Pauli Ch, Balling G, Lorenz HP, Eicken A, Hennig M, Hess J.](#)

[Cardiac index monitoring by pulse contour analysis and thermodilution after pediatric cardiac surgery](#)
[J of Thorac Cardiovasc Surg 2007; 133\(1\):224.8](#)

★★★

[Lopez–Herce J, Ruperez M, Sanchez C, Garcia C, Garcia E.](#)

[Estimation of the parameters of cardiac function and of blood volume by arterial thermodilution in an infant animal model](#)

[Pediatric Anesthesia 2006; 16: 635–40](#)

★★

[Lopez–Herce J, Ruperez M, Sanchez C, Garcia C, Garcia E.](#)

[Hemodynamic response to acute hypovolaemia, rapid blood volume expansion and adrenaline administration in an infant animal model](#)

[Resuscitation 2006; 68: 259–65](#)

★★

Egan J, Festa M, Cole A, Nunn GR, Gillis J, Winlaw DS.

Clinical assessment of cardiac performance in infants and children following cardiac surgery
Intensive Care Med 2005; 31(4):568-73

★★★

Torgay A, Pirat A, Akpek E, Zeyneloglu P, Arslan G, Haberal M.

Pulse contour cardiac output system use in pediatric orthotopic liver transplantation: preliminary report
of nine patients
Transplant Proc 2005; 37(7):3168-70

★★

Ruperez M, Lopez-Herce J, Garcia C, Sanchez C, Garcia E, Vigil D.

Comparison between cardiac output measured by the pulmonary arterial thermodilution technique and
that measured by the femoral arterial thermodilution technique in a pediatric animal model
Pediatr Cardiol 2004; 25(2):119-23

★★

Cecchetti C, Stoppa F, Vanacore N, Barbieri MA, Raucci U, Pasotti E, Tomasello C, Marano M, Pirozzi N.

Monitoring of intrathoracic volemia and cardiac output in critically ill children
Minerva Anesthesiol 2003; 69:907-18

★★★

Pauli C, Fakler U, Genz T, Hennig M, Lorenz HP, Hess J.

Cardiac output determination in children: equivalence of the transpulmonary thermodilution method to
the direct Fick principle
Intensive Care Med 2002; 28(7):947-52

★★★

Schiffmann H, Erdlenbruch B, Singer D, Singer S, Herting E, Hoefl A, Buhre W.

Assessment of cardiac output, intravascular volume status, and extravascular lung water by
transpulmonary indicator dilution in critically ill neonates and infants
J Cardiothorac Vasc Anesth 2002; 16(5):592-7

★★★

4.8 敗血症ショック(SHOCK SEPSIS)

Osman D, Ridel C, Ray P, Monnet X, Anguel N, Richard C, Teboul JL.

Cardiac filling pressures are not appropriate to predict hemodynamic response to volume challenge
Crit Care Med 2007; 35(1):64-9

★★★

[Kortgen A, Niederprun P, Bauer M.](#)

[Implementation of an evidence-based standard operating procedure“ and outcome in septic shock
Crit Care Med 2006; 34 \(4\):939-9](#)

★★★

[Martin GS, Eaton S, Mealer M, Moss M.](#)

[Extravascular lung water in patients with severe sepsis: a prospective cohort study
Crit Care 2005; 9\(2\):R74-82](#)

★★★

[Michard F, Alaya S, Zarka V, Bahloul M, Richard C, Teboul JL.](#)

[Global end-diastolic volume as an indicator of cardiac preload in patients with septic shock
Chest 2003; 124\(5\):1900-8](#)

★★★

4.9 出血性ショック(SHOCK HYPOVOLEMIA)

[Nirmalan M, Niranjana M, Willard T, Edwards JD, Little RA, Dark PM.](#)

[Estimation of errors in determining intrathoracic blood volume using thermal dilution in pigs with acute lung injury and haemorrhage
Br J Anaesth 2005; 93\(4\):546-51](#)

★★

[Berkenstadt H, Friedman Z, Preisman S, Keidan I, Livingstone D, Perel A.](#)

[Pulse pressure and stroke volume variations during severe haemorrhage in ventilated dogs
Br J Anaesth 2005; 94\(6\): 721-6](#)

★★

[Friedman Z, Berkenstadt H, Margalit N, Segal E, Perel A.](#)

[Cardiac output assessed by arterial thermodilution during exsanguination and fluid resuscitation: experimental validation against a reference technique
Eur J Anaesthesiol 2002; 19\(5\):337-40](#)

★★

4.10 心原性ショック(SHOCK CARDIOGENIC)

[Janda M, Scheeren TWL, Bajorat J, Westphal B, Vagts DA, Pohl B, Popescu C, Hofmockel R.](#)

[The impact of Intra-aortic Balloon Pumping on Cardiac Output Determination by Pulmonary Arterial and Transpulmonary Thermodilution in Pigs
J of Cardiovasc and Vasc Anesth 2006; 20 \(3\):320-4](#)

★★

[Bindels AJ, van der Hoeven JG, Graafland AD, de Koning J, Meinders AE.](#)

[Relationships between volume and pressure measurements and stroke volume in critically ill patients](#)

[Crit Care 2000; 4\(3\):193-9](#)

★

4.11 ARDS と ALI (ACUTE RESPIRATORY DISTRESS SYNDROME / ACUTE LUNG INJURY)

[Phillips, C., Chesnutt M, Smith, M.](#)

[Extravascular lung water in sepsis-associated acute respiratory distress syndrome: indexing with predicted body weight improves correlation with severity of illness and survival](#)

[Crit Care Med, 2007\(epub\): 1-5](#)

★★★

[Monnet X, Anguel N, Osman D, Hamzaoui, Richard C, Teboul JL.](#)

[Assessing pulmonary permeability by transpulmonary thermodilution allows differentiation of hydrostatic pulmonary edema from ALI / ARDS](#)

[Intensive Care Medicine 2007; 33\(3\): 448-53](#)

★★★

[Kuzkov V, Kirov M, Sovershaev M, Kulkin V, Suberov E, Waerhaug K, Bjertnaes L.](#)

[Extravascular lung water determined with single transpulmonary thermodilution correlates with the severity of sepsis-induced acute lung injury](#)

[Crit Care Med 2006; 34\(6\):1647-53](#)

★★★

[Perkins GD, McAuley DF, Thickett DR, Gao F.](#)

[The beta-agonist lung injury trial \(BALTI\): a randomized placebo-controlled clinical trial](#)

[Am J Respir Crit Care Med 2006; 173\(3\): 281-7](#)

★★★

[Martin GS, Eaton S, Mealer M, Moss M.](#)

[Extravascular lung water in patients with severe sepsis: a prospective cohort study](#)

[Crit Care 2005; 9\(2\):R74-82](#)

★★★

[Patroniti N, Bellani G, Maggioni E, Manfio A, Marcora B, Pesenti A.](#)

[Measurement of pulmonary edema in patients with acute respiratory distress syndrome](#)

[Crit Care Med 2005; 33\(11\):2547-54](#)

★★★

[McAuley DF, Giles S, Fichter H, Perkins GD, Gao F.](#)

[What is the optimal duration of ventilation in the prone position in acute lung injury and acute respiratory distress syndrome?](#)

[Intensive Care Med 2002; 28\(4\):414-8](#)

★

4.12 その他 (OTHERS)

脳神経外科関連 (NEUROSURGERY / NEUROLOGY)

[Mutoh T, Kazumata K, Ajiki M, Ushikoshi S, Terasaka S.](#)

[Goal-Directed Fluid Management by Bedside Transpulmonary Hemodynamic Monitoring After Subarachnoid Hemorrhage](#)

[Stroke, 2007 epub](#)

★★★

[Segal E, Greenlee JD, Hata SJ, Perel A.](#)

[Monitoring intravascular volumes to direct hypertensive, hypervolemic therapy in a patient with vasospasm](#)

[J Neurosurg Anesthesiol 2004; 16\(4\):296-8](#)

★★

[Berkenstadt H, Margalit N, Hadani M, Friedman Z, Segal E, Villa Y, Perel A.](#)

[Stroke volume variation as a predictor of fluid responsiveness in patients undergoing brain surgery](#)

[Anesth Analg 2001; 92\(4\):984-9](#)

★★

右-左シャント (RIGHT-LEFT SHUNT)

[Michard F, Alaya S, Medkour F.](#)

[Monitoring right-to-left intracardiac shunt in acute respiratory distress syndrome](#)

[Crit Care Med 2004; 32\(1\):308-9](#)

★★

肺水腫 (PULMONARY EDEMA)

[Groeneveld AB, Verheij J, van den Berg FG, Wisselink W, Rauwerda JA.](#)

[Increased pulmonary capillary permeability and extravascular lung water after major vascular surgery: effect on radiography and ventilatory variables](#)

[Eur J Anaesthesiol 2006; 23\(1\):36-41](#)

★★★

[Michard F, Schachtrupp A, Toens C.](#)

[Factors influencing the estimation of extravascular lung water by transpulmonary thermodilution in critically ill patients](#)

[Crit Care Med 2005; 33\(6\):1243-7](#)

★★★

[Patroniti N, Bellani G, Maggioni E, Manfio A, Marcora B, Pesenti A.](#)

[Measurement of pulmonary edema in patients with acute respiratory distress syndrome](#)

[Crit Care Med 2005; 33\(11\):2547–54](#)

★★★

[Boussat S, Jacques T, Levy B, Laurent E, Gache A, Capellier G, Neidhardt A.](#)

[Intravascular volume monitoring and extravascular lung water in septic patients with pulmonary edema](#)

[Intensive Care Med 2002; 28\(6\):712–8](#)

★★★

5. 各パラメータによる治療補助(DECISION SUPPORT BY PICCO PARAMETERS)

[Goepfert M, Reuter D, Akyol D, Lamm P, Kilger E, Goetz A.](#)

[Goal directed fluid management reduces vasopressor and catecholamine use in cardiac surgery patients](#)

[Intensive Care Medicine 2007; 33: 96–103](#)

★★★

[Compton F, Hoffmann C, Zidek W, Schmidt S, Schaefer JH.](#)

[Volumetric hemodynamic parameters to guide fluid removal on hemodialysis in the intensive care unit](#)

[Hemodial Int 2007; 11\(2\):231–7](#)

★★★

[Hensel M, Schwenk W, Bloch A, Raue W, Stracke S, Volk T, v. Heymann C, Muller JM, Kox WJ, Spies C.](#)

[\[Die Aufgabe der Anesthesiologie bei der Umsetzung operativer „Fast track-Konzepte“\] \(Article in German\)](#)

[Der Anesthesist 2006; 5\(1\):80–92](#)

★★

[Holm C, Tegeler J, Mayr M, Pfeiffer U, Henckel von Donnersmarck G, Muhlbauer W.](#)

[Effect of crystalloid resuscitation and inhalation injury on extravascular lung water: clinical implications](#)

[Chest 2002; 121\(6\):1956–62](#)

★★

[Holm C, Melcer B, Horbrand F, Worl H, von Donnersmarck GH, Muhlbauer W.](#)

[Intrathoracic blood volume as an end point in resuscitation of the severely burned: an observational study of 24 patients](#)

[J Trauma 2000; 48\(4\):728–34](#)

★★★

[Mitchell JP, Schuller D, Calandrino FS, Schuster DP.](#)

[Improved outcome based on fluid management in critically ill patients requiring pulmonary artery catheterization](#)

[Am Rev Respir Dis 1992; 145\(5\):990–8](#)

★★★

6. 様々な臨床状態におけるパラメータ(PICCO PARAMETERS IN VARIOUS CLINICAL SITUATIONS)

[Sakka S, Hanusch T, Thuemer , Wegscheider K.](#)

[The influence of venovenous renal replacement therapy on measurements by the transpulmonary thermodilution technique](#)

[Anesth Analg 2007; 105\(4\):1079-82](#)

★★★

[Michelet P, Roch A, Gainnier M, Sainty JM, Auffray JP, Papazian L.](#)

[Influence of support on intra-abdominal pressure, hepatic kinetics of indocyanine green and extravascular lung water during prone positioning in patients with ARDS: a randomized crossover study](#)

[Crit Care 2005; 9\(3\):R251-7](#)

★★

[Wiesenack C, Prasser C, Liebold A, Schmid FX.](#)

[Assessment of left ventricular cardiac output by arterial thermodilution technique via a left atrial catheter in a patient on a right ventricular assist device](#)

[Perfusion 2004; 19\(1\):73-5](#)

★

[Luecke T, Roth H, Herrmann P, Joachim A, Weisser G, Pelosi P, Quintel M.](#)

[PEEP decreases atelectasis and extravascular lung water but not lung tissue volume in surfactant washout lung injury](#)

[Intensive Care Med 2003; 29\(11\):2026-33](#)

★★

[Reuter DA, Felbinger TW, Schmidt C, Moerstedt K, Kilger E, Lamm P, Goetz AE.](#)

[Trendelenburg positioning after cardiac surgery: effects on intrathoracic blood volume index and cardiac performance](#)

[Eur J Anaesthesiol 2003; 20\(1\):17-20](#)

★★

[Hofer CK, Zalunardo MP, Klaghofer R, Spahr T, Pasch T, Zollinger A.](#)

[Changes in intrathoracic blood volume associated with pneumoperitoneum and positioning](#)

[Acta Anaesthesiol Scand 2002; 46\(3\):303-8](#)

★★

[Kampen J, Liess K, Casadio C, Tonner PH, Scholz J.](#)

[\[Thermal lesions caused by a PiCCO catheter left in place in the MRT\] \(Article in German\)](#)

[Intensivmedizin und Notfallmedizin 2002; 39\(1\):113](#)

★★

[Buhre W, Weyland A, Buhre K, Kazmaier S, Mursch K, Schmidt M, Sydow M, Sonntag H.
Effects of the sitting position on the distribution of blood volume in patients undergoing neurosurgical
procedures](#)

[Br J Anaesth 2000; 84\(3\):354-7](#)

★★

7. カテーテル関連(CATHETERS)

[Schmidt S, Westhoff TH, Hofmann C, Schaefer JH, Zidek W, Compton F, van der Giet M.
Effect of the venous catheter site on transpulmonary thermodilution measurement variables](#)

[Crit Care Med 2007; 35\(3\):783-6](#)

★★★

[Traore O, Liotier J, Souweine B.](#)

[Prospective study of arterial and central venous catheter colonization and of arterial- and central venous
catheter-related bacteremia in intensive care units](#)

[Crit Care Med 2005; 33\(6\):1276-80](#)

★★★

[Cousins TS, O'Connell JM.](#)

[Arterial Cannulation: A Critical Review](#)

[AANA Journal 2004; 72 \(4\): 267-71](#)

★★★

[Orme RM, Pigott DW, Mihm FG.](#)

[Measurement of cardiac output by transpulmonary arterial thermodilution using a long radial artery
catheter. A comparison with intermittent pulmonary artery thermodilution](#)

[Anaesthesia 2004; 59: 590-4](#)

★★★

[Segal E, Katzenelson R, Berkenstadt H, Perel A.](#)

[Transpulmonary thermodilution cardiac output measurement using the axillary artery in critically ill
patients](#)

[J Clin Anesth 2002; 14\(3\):210-3](#)

★★★

[Scheer B, Perel A, Pfeiffer UJ.](#)

[Clinical review: complications and risk factors of peripheral arterial catheters used for haemodynamic
monitoring in anaesthesia and intensive care medicine](#)

[Crit Care 2002; 6\(3\):199-204](#)

★★★

Clementi G.

Hemodynamic monitoring using a long radial catheter

Minerva Anesthesiol 2002; 68: 231 – 5

